arXiv:hep-ph/0501255v2 30 Sep 2005

Large Angle Hadron Correlations from Medium-Induced Gluon Radiation*

Ivan Vitev
Los Alamos National Laboratory, Theory Division and Physics Division, Mail Stop H846, Los Alamos, NM 87545, USA

(Dated: February 2, 2008)

Final state medium-induced gluon radiation in ultradense nuclear matter is examined and shown to favor large angle emission when compared to vacuum bremsstrahlung due to the suppression of collinear gluons. Perturbative expression for the contribution of its hadronic fragments to the back-to-back particle correlations is derived. It is found that in the limit of large jet energy loss gluon radiation determines the yield and angular distribution of $|\Delta\varphi| \geq \frac{\pi}{2}$ di-hadrons to high transverse momenta p_{T_2} of the associated particles. Clear transition from enhancement to suppression of the away-side hadron correlations is established at moderate p_{T_2} and its experimentally accessible features are predicted versus the trigger particle momentum p_{T_1}.

PACS numbers: 12.38.Bx,12.38.Mh,25.75.-q,25.75.Gz

The discovery of jet quenching [1] – the suppression of large transverse momentum hadron production in nuclear collisions relative to the expectation from p+p reactions scaled by the number of elementary nucleon-nucleon interactions – is arguably the most exciting new result from the Au+Au experimental program at the Relativistic Heavy Ion Collider (RHIC). The phenomenon of jet quenching has been established via the attenuation of the single inclusive particle spectra [2] and the suppression of the back-to-back di-hadron correlations [3, 4]. It has been interpreted as critical evidence for large parton energy loss [5, 6, 7] in ultradense quark-gluon plasma (QGP) – the deconfined state of matter predicted by quantum chromodynamics (QCD).

So far, the observable effects of the medium-induced gluon radiation *itself* were considered to be modest. For tagged jets, assuming thermalization of the lost energy calculated in [8], transport models predicted an increase in the multiplicity of associated particles limited to $p_{T_2} \leq 500$ MeV [9]. Using the angular gluon distribution from [10], jet cone broadening was found to be $< 10\%$ and challenging to detect experimentally even at the CERN Large Hadron Collider (LHC) [11].

In this Letter we demonstrate that a mechanism, based on the destructive interference of color currents from hard and soft parton scattering, can ensure a broad (in angle and frequency) final state medium-induced emission spectrum in QCD. For large energy loss, perturbative fragmentation of the radiative gluons is found to give a dominant contribution to the yield of away-side di-hadrons and to significantly alter their correlations at transverse momenta much higher than naively anticipated.

We first recall that for hard perturbative scattering the radiative spectrum of real gluon emission for small and moderate frequencies ω is given by [8]

$$\frac{dN^g_{\rm vac}}{d\omega d\sin\theta^* d\delta} \approx \frac{C_R\alpha_s}{\pi^2}\frac{1}{\omega\sin\theta^*} \, . \qquad (1)$$

In Eq. (1) $\theta^* = \arcsin(k_\perp/\omega)$ is the angle relative to the jet axis, δ is the azimuthal cone angle (both illustrated in Fig. 1), α_s is the strong coupling constant and $C_R = 3\,(4/3)$ for gluon (quark) jets, respectively. Virtual gluon corrections remove the $\omega \to 0$ infrared singularity in the cross sections in accord with the Kinoshita-Lee-Nauenberg theorem [12] but the collinear $\theta^* \to 0$ divergence has to be regulated or subtracted in the parton distribution functions (PDFs) and the fragmentation functions (FFs).

In contrast, the final state medium-induced bremsstrahlung spectrum is both collinear and infrared safe. To first order in the mean number of soft interactions in the plasma the Gyulassy-Levai-Vitev [8] gluon distribution in angle and frequency reads:

$$\frac{dN^g_{\rm med}}{d\omega d\sin\theta^* d\delta} = \frac{2C_R\alpha_s}{\pi^2}\int_{z_0}^{L}\frac{d\Delta z}{\lambda_g(z)}\int_0^\infty dq_\perp\, q_\perp^2\,\frac{1}{\sigma_{el}}\frac{d\sigma_{el}}{d^2q_\perp}(z)$$
$$\times\int_0^{2\pi} d\alpha\,\frac{\cos\alpha}{q_\perp^2 - 2q_\perp\omega\sin\theta^*\cos\alpha + \omega^2\sin^2\theta^*}$$
$$\times\left[1-\cos\left(\frac{(q_\perp^2 - 2q_\perp\omega\sin\theta^*\cos\alpha + \omega^2\sin^2\theta^*)\Delta z}{2\omega}\right)\right] \, . \qquad (2)$$

In Eq. (2) $\alpha = \angle(\vec{k}_\perp, \vec{q}_\perp)$, $\lambda_g(z)$ is the position-dependent gluon mean free path and L is the transverse size of the medium. The momentum transfers $\vec{q}_\perp$ are distributed according to the normalized elastic scattering cross section $\sigma_{el}^{-1} d\sigma_{el}(z)/d^2q_\perp = \mu_D^2(z)\pi^{-1}(q_\perp^2+\mu_D^2(z))^{-2}$. In this model, $\langle q_\perp^2\rangle \propto \mu_D^2(z)$ and for a quark-gluon plasma in local thermal equilibrium $\mu_D^2(z) \sim 4\pi\alpha_s T^2$. For the case of (1+1)D dynamical Bjorken expansion of the QGP $\mu_D(z) = \mu_D(z_0)\,(z_0/z)^{1/3}$, $\lambda_g(z) = \lambda_g(z_0)\,(z/z_0)^{1/3}$[5, 6].

---

[*] On September 25, 2005 my advisor, friend and colleague Ventseslav Rizov passed away. With his untimely death the Bulgarian physics community lost a person of great kindness, integrity and devotion to science. This work is dedicated to his memory.



FIG. 1: The angular distribution of medium-induced bremsstrahlung of $E = 6$ GeV gluon jet for fixed values of the radiative gluon energy $\omega = 0.5, 1, 2, 4$ GeV. Top and bottom panels represent (1+1)D Bjorken expanding medium of transverse size $L = 9$ fm and $L = 3$ fm, respectively. Inserts show $\langle\theta^*\rangle$ versus ω. Right panels illustrate the gluon spectrum and radiation intensity of $E = 40$ GeV gluon jet.

From Eq. (2) the gluon distribution is not only finite when $\theta^* \to 0$ but vanishes on average due to the uniform angular distribution of momentum transfers from the medium, $\int_0^{2\pi} d\alpha \cos\alpha = 0$. We have checked that for physical gluons of $k_\perp \leq \omega$ the cancellations discussed here persist to all orders in the mean number of scatterings [8]. The small frequency and small angle spectral behavior of $dN^g_{\rm med}/d\omega d\sin\theta^* d\delta$ remains under perturbative control. We also emphasize that destructive quantum interference suppresses radiation of $\Delta z \ll l_f$ [8]. The induced radiation decouples from the jet at a scale $l_f \sim \Delta z \propto L/2$ and facilitates hadronization outside of the medium.

Given the vastly different angular behavior of the vacuum and the medium-induced gluon bremsstrahlung, Eqs. (1) and (2), it is critical to identify the phase space where cancellation of the color currents induced by the hard and soft scattering occurs. We fix the parameters of the medium in Eq. (2) to $\mu_D(z_0) = 1.5$ GeV and $\lambda_g(z_0) = 0.75$ fm at initial formation time $z_0 = 0.25$ fm. Since small frequency emission is suppressed, we use only a moderate $\alpha_s = 0.25$. Triggering on high p_{T_1} hadron directs its parent parton "c" away from the medium and places the collision point of the lowest order (LO) ab → cd underlying perturbative process [13] close to the periphery of the nuclear overlap region. Then, it is the back-scattered jet "d" that traverses the QGP. For large nuclei, such as Au and Pb, path lengths $L = 9$ (3) fm are used to illustrate central (peripheral) collisions, respectively. We limit gluon emission to the forward jet hemisphere, $0 \leq \theta^* \leq \frac{\pi}{2}$.

The angular distribution of medium-induced radiation for $E = 6$ GeV gluon jet for select values of ω is shown in the right panels of Fig. 1. We find that gluon emission is strongly suppressed within a cone of opening angle $\theta^* \simeq 0.25$ rad due to the cancellation of collinear bremsstrahlung – a mechanism different from a Gaussian random walk in θ^*. The broad gluon distribution can be characterized by the mean emission angle

$$\langle\theta^*\rangle = \int_0^1 \theta^* \frac{dN^g_{\rm med}}{d\omega d\sin\theta^*} \, d\sin\theta^* \left[\int_0^1 \frac{dN^g_{\rm med}}{d\omega d\sin\theta^*} \, d\sin\theta^* \right]^{-1},$$

given in the inserts of Fig. 1. Left panels illustrate the infrared safety property of the spectrum. Energy loss is dominated by semihard few GeV gluons, see $\omega dN^g/d\omega$, emitted at $\langle\theta^*\rangle \sim 0.5$ rad.

Results reported in this Letter are important for the future LHC heavy ion program since they for the first time suggest a large, and therefore detectable, broadening of the jet cone in the nuclear environment. Note that $\langle\theta^*\rangle$ implies a redistribution of the energy flow even within jets of large opening angle $R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 1$. At present, however, a key question for perturbative QCD phenomenology is whether the medium induced gluon bremsstrahlung can significantly alter the di-hadron correlations measured at RHIC [3, 4, 14]. We naturally focus on the away-side $|\Delta\varphi| \geq \frac{\pi}{2}$ case, where medium effects are the largest. Nuclear modifications build upon the LO double inclusive hadron production cross section, which is calculable in the perturbative QCD factorization approach [15] if either of the hadrons is moderately hard (p_{T_1} or $p_{T_2} \geq$ few GeV) [13, 16]:

$$\frac{d\sigma^{h_1h_2}_{NN}}{dy_1 dy_2 dp_{T_1} dp_{T_2} d\Delta\varphi} = K \sum_{abcd} \int_0^1 \frac{dz_1}{z_1} D_{h_1/c}(z_1) \times \left[D_{h_2/d}(z_2) \delta(\Delta\varphi - \pi) \right] \frac{\phi_{a/N}(x_a)\phi_{b/N}(x_b)}{x_a x_b \, S^2} \times 2\pi\alpha_s^2 |\overline{M}_{ab\to cd}|^2 \, . \quad (3)$$

In Eq. (3) $K = 2$ is a next-to-leading order K-factor, $x_{a,b} = p_{a,b}/p_{N_a,N_b}$ are the momentum fractions of the incoming partons and $z_{1,2} = p_{h_1,h_2}/p_{c,d}$ are the momentum fractions of the hadronic fragments. We use standard lowest order Gluck-Reya-Vogt PDFs [17] and Binnewies-Kniehl-Kramer FFs [18]. Renormalization, factorization and fragmentation scales are suppressed everywhere for clarity. The spin (polarization) and color averaged matrix elements $|\overline{M}_{ab\to cd}|^2$ are given in [13].

We shall first discuss the physical effects that alter the LO perturbative formula, Eq. (3). A modification that does not change the $\Delta\varphi$-integrated cross section is vacuum- and medium-induced acoplanarity. The deviation of jets from being back-to-back in a plane perpendicular to the collision axis arises from the soft gluon radiation and transverse momentum diffusion in dense



FIG. 2: Nuclear modification of the back-to-back di-hadron correlations with and without the contribution of medium-induced bremsstrahlung. Top and bottom panels illustrate central and peripheral collisions, respectively. Data is from STAR [4]. Right panels predict $R^{(2)}_{AA}$ for low and high p_{T_1} hadron triggers.

nuclear matter [19]. In the approximation of collinear fragmentation, the width of the away-side hadron-hadron correlation function can be related to the accumulated di-jet transverse momentum squared in the φ-plane, $\sin\sqrt{\frac{2}{\pi}}\sigma_{\rm Far} = \sqrt{\frac{2}{\pi}\langle k_T^2\rangle_\varphi}/p_{\perp_d}$. Assuming a Gaussian form,

$$f_{\rm vac.\ or\ med.}(\Delta\varphi) = (\sqrt{2\pi}\sigma_{\rm Far})^{-1}\exp[-(\Delta\varphi-\pi)^2/2\sigma^2_{\rm Far}] ,$$

a good description of $|\Delta\varphi| \geq \frac{\pi}{2}$ correlations measured in elementary p+p collisions at $\sqrt{S} = 200$ GeV [7] requires a large $\langle k^2_{T\ \rm vac}\rangle_\varphi = 5\ {\rm GeV}^2$ for the di-jet pair with away-side scattered quark (and a 2.25 larger value for a scattered gluon). Additional broadening arises from the interactions of the jet in the QGP that ultimately lead to the reported energy loss. Using the parameters of the medium for (1+1)D expansion we find

$$\langle k^2_{T\,\rm hot}\rangle = \int_{z_0}^{L} dz\ 2\frac{\mu^2_D(z)}{\lambda_{q,g}(z)} = 2\frac{\mu^2_D(z_0)}{\lambda_{q,g}(z_0)}\ln\frac{L}{z_0} , \qquad (4)$$

although only half is projected on the φ-plane, $\langle k^2_T\rangle_\varphi = \langle k^2_{T\ \rm vac}\rangle_\varphi + \frac{1}{2}\langle k^2_{T\ \rm hot}\rangle$.

Two competing mechanisms do, however, change the p_{T_2} dependence of the perturbative cross section, Eq. (3). First is the the parent jet "d" fractional energy loss $\epsilon = \Delta E_d/E_d$, which we here for simplicity consider on average and evaluate by integrating Eq. (2). It leads to a rescaling of the hadronic fragmentation momentum fraction $z_2 \to z_2/(1-\epsilon)$ [5]. Physically, fewer high p_{T_2} particles are produced by the attenuated parton of energy $E_d - \Delta E_d$. If the energy loss is large, a second mechanism is invoked as a consequence. Hadronic fragments of the radiative gluons will increase the probability of finding low and moderate p_{T_2} particles associated with the interacting jet [9].

To calculate di-hadron correlations, we first map the jet structure of a hard 90^0- scattered parton on rapidity $y \approx \eta = -\ln\tan(\theta/2)$ (θ being the angle relative to the beam axis) and azimuth ϕ, $\tan^2\theta^* = \cot^2\theta + \tan^2\phi$, $\tan\delta = -\cot\theta/\tan\phi$. The approximately flat rapidity distribution of the away-side jet near $y_2 = 0$ can be used to sum over all emission angles $\theta \in (\theta_{\rm min}, \theta_{\rm max}) \subset (0,\pi)$ yielding

$$\frac{dN^g_{\rm med}}{d\omega d\phi} = \int_{\theta_{\rm min}}^{\theta_{\rm max}} d\theta\ \left[\frac{dN^g_{\rm med}}{d\omega d\sin\theta^* d\delta}\left|\frac{\partial(\sin\theta^*,\delta)}{\partial(\theta,\phi)}\right|\right] . \qquad (5)$$

The Jacobian of the transformation in Eq. (5) reads

$$\left|\frac{\partial(\sin\theta^*,\delta)}{\partial(\theta,\phi)}\right| = \frac{1}{\sin^2\theta\cos^2\phi}\frac{(\tan^2\phi+\cot^2\theta)^{-1/2}}{(1+\tan^2\phi+\cot^2\theta)^{3/2}} .$$

It is critical to note that projection on a plane coincident with the jet cone axis (the ϕ-plane in Eq. (5) is one example) efficiently masks the $\theta^* \to 0$ void in the angular distribution of medium-induced gluons reported in Fig. 1. Our conclusion is independent of the physical mechanism that depletes the parton (or particle) multiplicity in a cone around the jet axis and important for the interpretation of the experimental data.

The end analytic result for the modification to Eq. (3) per average nucleon-nucleon collision in the heavy ion environment can be derived from the energy sum rule for all hadronic fragments from the jet,

$$\begin{aligned} & D_{h_2/d}(z_2)\delta(\Delta\varphi-\pi) \quad \Rightarrow \\ & \frac{1}{1-\epsilon}D_{h_2/d}\left(\frac{z_2}{1-\epsilon}\right) f_{\rm med.}(\Delta\varphi) + \frac{p_{T_1}}{z_1}\int_0^1\frac{dz_g}{z_g}D_{h_2/g}(z_g) \\ & \qquad \times\int_{-\pi/2}^{\pi/2}d\phi\ \frac{dN^g_{\rm med}(\phi)}{d\omega d\phi}f_{\rm vac.}(\Delta\varphi-\phi) . \qquad (6)\end{aligned}$$

Here, $z_g = p_{T_2}/\omega$.

Whether medium-induced gluon radiation may have significant observable consequences for the large angle di-hadron correlations depends on its relative contribution to the $\Delta\varphi$-integrated cross section. From Eq. (3) this can be studied versus p_{T_2} via the ratio [16]

$$R^{(2)}_{AA} = \frac{d\sigma^{h_1h_2}_{AA}/dy_1dy_2dp_{T_1}dp_{T_2}}{\langle N^{\rm coll}_{AA}\rangle\, d\sigma^{h_1h_2}_{NN}/dy_1dy_2dp_{T_1}dp_{T_2}} , \qquad (7)$$

where the mean number of collisions $\langle N^{\rm coll}_{AA}\rangle$ is estimated from an optical Glauber model [2, 3]. Numerical results, shown in Fig. 2, correspond to triggering on a high $p_{T_1} = 4$ - 6 GeV pion and measuring all associated $\pi^+ + \pi^0 + \pi^-$. Depletion of hadrons from the quenched parent parton alone leads to a large suppression of the



FIG. 3: The angular distribution of $|\Delta\varphi| \geq \frac{\pi}{2}$ di-hadrons for central and peripheral A+A collisions. Solid and dashed lines give *separately* the contribution of the radiative gluons and the attenuated parent parton. The lower right panel compares the the width of the correlation function in p+p to the one of the suppressed parent parton in central A+A reactions.

double inclusive cross section with weak p_{T_2} dependence. Hadronic feedback from the medium-induced gluon radiation, however, completely changes the nuclear modification factor $R^{(2)}_{AA}$. It now shows a clear transition from a quenching of the away-side jet at high transverse momenta to enhancement at $p_{T_2} \leq 2$ GeV, a scale significantly larger than the one found in [9]. In fact, we have checked that for large ΔE_d the back-to-back di-hadron correlations are dominated by radiative gluons to unexpectedly high $p_{T_2} \sim 10$ GeV. STAR data in central and peripheral Au+Au collisions [4] is shown for comparison. Experimentally testable predictions for the shape and magnitude of $R^{(2)}_{AA}$ and the enhancement-to-suppression transition $R^{(2)}_{AA} = 1$ versus p_{T_1} are also given in Fig. 2. It has been previously argued [20] that detailed balance reduce jet suppression at partonic scales $p_{\perp_d} \leq$ few μ_D, though such mechanism cannot produce enhancement of the low transverse momentum particle production observed in Fig. 2.

With a dominant contribution to the di-hadron yields, the medium induced gluons are bound to also determine the width of the correlation function. Two-particle distributions in A+A reactions, calculated from Eqs. (3) and (6) for a $p_{T_1} = 6$ GeV trigger pion and two different $p_{T_2} = 1,\ 4$ GeV associated pions, are shown in Fig. (3). Qualitatively, the medium-induced gluon component to the cross section controls the growth of the correlation width in central and semi-central nuclear collisions. Quantitatively, the effect should be even larger than the one estimated here, which is limited by the imposed $0 < \theta^* < \frac{\pi}{2}$ constraint. The decisive role of the bremsstrahlung spectrum, Eq. (2), in establishing the $\Delta\varphi$-shape of away-side di-hadrons is further clarified in the bottom right panel of Fig. 3. Hadronic fragments from the quenched parton "d" *only* are shown to yield a distribution that is *not* broader than the one anticipated in p+p reactions. Experimental measurements of significantly enhanced widths for $|\Delta\varphi| \geq \frac{\pi}{2}$ two-particle correlation in A+A collisions should thus point to copious hadron production from medium-induced large angle gluon emission.

In summary, we calculated the transverse momentum and angular distribution of the away-side di-hadron correlations in the framework of the perturbative QCD factorization approach, augmented by inelastic jet interactions in the quark-gluon plasma. At RHIC energies we found that the medium-induced gluon radiation determines the two-particle yields and the width of their correlation function to surprisingly high transverse momentum $p_{T_2} \sim 10$ GeV. Clear transition from back-to-back jet enhancement to back-to-back jet quenching is established at moderate $p_{T_2} = 1 - 3$ GeV, independent of collision centrality but sensitive to the trigger hadron momentum p_{T_1}. Definitive experimental determination of its features, predicted in this Letter, will for the first time provide a handle on the *differential spectrum* of medium-induced non-Abelian bremsstrahlung. Coincidental confirmation of large broadening of the away-side di-hadrons would require a critical reassessment of the origin of intermediate transverse momentum particles in central and semi-central nuclear collisions. For jet physics studies at the LHC, our findings ensure a measurable increase in the jet width in ultradense nuclear matter.

Helpful discussion with P. Constantin, B. Jacak, M. Johnson and F. Wang is gratefully acknowledged. This work is supported by the J. R. Oppenheimer Fellowship of the Los Alamos National Laboratory and by the US Department of Energy.

[1] P. Levai *et al.*, Nucl. Phys. A **698**, 631 (2002); K. Adcox *et al.*, Phys. Rev. Lett. **88**, 022301 (2002).
[2] S. S. Adler *et al.*, Phys. Rev. Lett. **91**, 072301 (2003).
[3] C. Adler *et al.*, Phys. Rev. Lett. **90**, 082302 (2003).
[4] J. Adams *et al.*, nucl-ex/0501016; F. Wang, J. Phys. G **30**, S1299 (2004).
[5] M. Gyulassy, I. Vitev, X. N. Wang, Phys. Rev. Lett. **86**, 2537 (2001). I. Vitev, M. Gyulassy, Phys. Rev. Lett. **89**, 252301 (2002).
[6] X. N. Wang, Phys. Lett. B **595**, 165 (2004).
[7] J. Adams *et al.*, Phys. Rev. Lett. **91**, 072304 (2003).
[8] M. Gyulassy, P. Levai, I. Vitev, Phys. Rev. Lett. **85**, 5535 (2000); Nucl. Phys. B **594**, 371 (2001).
[9] S. Pal, S. Pratt, Phys. Lett. B **574**, 21 (2003).
[10] R. Baier *et al.*, Phys. Rev. C **60**, 064902 (1999).
[11] C. A. Salgado, U. A. Wiedemann, Phys. Rev. Lett. **93**, 042301 (2004).
[12] T. Kinoshita, J. Math. Phys. **3**, 650 (1962); T. D. Lee, M. Nauenberg, Phys. Rev. **133**, B1549 (1964).

[13] J. F. Owens, Rev. Mod. Phys. **59**, 465 (1987).

[14] J. Rak, J. Phys. G **30**, S1309 (2004).

[15] J. C. Collins, D. E. Soper, G. Sterman, Nucl. Phys. B **261**, 104 (1985).

[16] J. W. Qiu, I. Vitev, hep-ph/0405068.

[17] M. Gluck, E. Reya, A. Vogt, Eur. Phys. J. C **5**, 461 (1998).

[18] J. Binnewies, B. A. Kniehl, G. Kramer, Z. Phys. C **65**, 471 (1995).

[19] J. W. Qiu, I. Vitev, Phys. Lett. B **570**, 161 (2003).

[20] E. Wang, X. N. Wang, Phys. Rev. Lett. **87**, 142301 (2001).